

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 7/48, 7/06	A1	(11) International Publication Number: WO 95/22311 (43) International Publication Date: 24 August 1995 (24.08.95)
(21) International Application Number: PCT/EP95/00582 (22) International Filing Date: 16 February 1995 (16.02.95) (30) Priority Data: 9403156.4 18 February 1994 (18.02.94) GB 9414332.8 15 July 1994 (15.07.94) GB (71) Applicant (for AU BB CA GB IE KE LK MN MW NZ SD SZ TT UG only): UNILEVER PLC [GB/GB]; Unilever House, Blackfriars, London EC4P 4BQ (GB). (71) Applicant (for all designated States except AU BB CA GB IE KE LK MN MW NZ SD SZ TT UG): UNILEVER NV [NL/NL]; Weena 455, NL-3013 AL Rotterdam (NL). (72) Inventors: HAGUE, Jonathon, David; 24 Trafalgar Drive, Bebington, Wirral, Merseyside L63 7RW (GB). MURRAY, Andrew, Malcolm; 15 Moorings Close, Parkgate, South Wirral, Cheshire L64 6TL (GB). (74) Agent: MULDER, Cornelis, W.; Unilever plc, Patent Division, Colworth House, Sharnbrook, Bedford MK44 1LQ (GB).		(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, UG, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: PERSONAL WASHING COMPOSITIONS (57) Abstract Personal washing compositions which contain a cationic deposition polymer of charge density in the range 0.0001 to 0.005 eq/g and average molecular weight greater than 2×10^6 daltons in combination with a surfactant to increase deposition of a benefit agent dispersed in the composition onto the skin or hair.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

PERSONAL WASHING COMPOSITIONS

This invention relates to personal washing compositions which contain a cationic deposition polymer to increase deposition of a benefit agent onto the skin or hair.

For many years it has been known that hair can be conditioned by incorporation of silicone in a shampoo composition. US 2 826 551 (Geen) is typical of an early disclosure of such 2 in 1 shampoos. Various attempts have been made to improve the efficiency of use of the expensive silicone component. This would provide better conditioning and the option of reducing the level of expensive benefit agent in the shampoo, with consequent cost saving.

Deposition polymers with a cationic charge have been proposed to enhance the amount of benefit agent deposited from the shampoo. For example cationic guar gum has been described for the enhancement of the deposition of antidandruff particles in US 5 037 818 and for the enhanced deposition of insoluble non-volatile silicone in US 5 085 857. The use of cationic polymers in shower gels to enhance deposition of silicone oil is also known from EP-A-457 688 (L'Oreal).

Deposition polymers have also been proposed to enhance the deposition of sunscreen materials from a shampoo composition. In EP 386 898 a cationic polygalactomannan gum derivative is used.

Polyacrylamides have been proposed for use in shampoos in EP 0 231 997. These polymers are not charged and do not assist in the deposition of benefit agents.

When washing with any of the prior art systems a considerable amount of the benefit agent will be rinsed away with the


- 2 -

composition, and there is scope for substantially improving the deposition efficiency.

5 It is an object of the present invention to provide a more efficient deposition polymer than the previously described polygalactomannan polymers.

10 According to the present invention there is provided a personal washing composition comprising: a surface active agent selected from anionic, nonionic, zwitterionic and cationic surfactants, soap and mixtures thereof, water, a non-volatile insoluble benefit agent dispersed in the composition and from 0.001 to 1% by weight of a deposition polymer which is a cationic copolymer wherein the charge
15 density of the copolymer is in the range 0.0001 to 0.005 eq/g, preferably 0.0008 to 0.0025 eq/g; and the average molecular weight of the copolymer is more than 2×10^6 daltons.

20 Preferably the amount of deposition polymer lies in the range 0.05 to 0.2% by weight. Preferably the cationic copolymer is a copolymer of acrylamide and a cationic monomer having the formula:



where: T is -O- or -C-



35 R is H or CH_3

and R' is $-\text{NH}-(\text{CH}_2)_n-\text{N}^+(\text{CH}_3)_3 \text{X}^-$ or $-\text{O}-(\text{CH}_2)_n-\text{N}^+(\text{CH}_3)_3 \text{X}^-$

- 3 -

in which n is an integer from 1 to 4 and X is selected from Cl, Br, I and CH_3SO_3 .

5 The composition is suitable for cleansing and conditioning of the skin or hair. The term "conditioning" is intended to cover "moisturising" and "protection".

Throughout this specification reference to average molecular weight (Mw) means a molecular weight calculated as follows.

10 The intrinsic viscosity of a polymer may be determined by standard capillary viscometry. The viscosity of a series of low polymer concentrations in a given solvent is determined relative to the pure solvent. The relative viscosity N_r is
15 defined as:

$$N_{\text{polymer sol}}/N_{\text{solvent}}$$

The specific viscosity N_{sp} is:

20
$$N_r - 1$$

If N_{sp}/c , where c is the polymer concentration, is plotted against c, a straight line is usually obtained. The point at
25 which the straight line crosses the y intercept is the intrinsic viscosity (N_{in}). This is related to the coil size of the polymer.

The intrinsic viscosity can also be related to the molecular
30 weight of the polymer if the Mark-Houwink parameters are known. Thus,

$$N_{in} = K(Mw)^a$$

35 Where K and a are the Mark-Houwink parameters.

- 4 -

These have been determined for polyacrylamide, and also for a number of copolymers of acrylamide and N,N,N trimethylaminoethyl chloride acrylate. These polymers are in accordance with those found useful in the present invention.

5 The parameters can be found in Mabire et al, Polymer 1984 (25) 1984.

For CPA1, CPA2 and CPA5 to 10 we used the MH parameters for 30% cationic polymer. For CPA 3 and CPA 4 we used MH

10 parameters for acrylamide homopolymer. Whichever is used, the Mw figure is approximately the same. Thus, if the MH parameters for acrylamide are used for CPA 1, the Mw figure becomes 5,000,000 rather than 8,000,000.

15 There is a precedent in the scientific literature where an estimate of Mw has been made for cationic polyacrylamides by using the MH parameters for acrylamide homopolymer: Hubbe, M.A, Colloids and Surfaces 1987 25 p. 325.

20 The MH parameters are generated for 1 M NaCl, so, in accordance with normal practice for polyelectrolytes, 1M NaCl was used as a solvent for our measurements. Intrinsic viscosity of some of the polymers used is given below:

- 5 -

Polymer	Intrinsic Viscosity (dl/g) 1M NaCl
CPA 1	11.1*
CPA 2	1.1*
CPA 3	11.8
CPA 4	1.2
CPA 5	12.5
CPA 6	8.0
CPA 7	2.8
CPA 9	10.4
CPA 9	6.9
CPA 10	8.25
JR 400	4.8
JR 30M	12.0
Jaguar C13S	9.8
Merquat 550	2.7

"CPA" polymers are copolymers of acrylamide and N,N,N-trimethyl aminopropylacrylamide. CPA 1, 2, 5, 7, 9 and 10 have cationic charge densities of 0.00145 eq/g. CPA 3 and CPA 4 have cationic charge densities of 0.0004 eq/g. CPA 6 and CPA 8 have respective cationic charge densities 0.00194 and 0.0009 eq/g. All CPA copolymers were ex Allied colloids. The Intrinsic viscosities for CPA 1, 2 and 5 to 10 is data from Allied Colloids. The data for CPA3 and CPA4 was produced by the above method and corresponds closely with the data supplied by Allied Colloids for these materials.

- 6 -

Preferred "CPA" polymers have an intrinsic viscosity of at least 7.

By "benefit agent" is meant a protective and/or softening substance that maintains softness by retarding the decrease in water content from the skin (stratum corneum) or hair.

Normally the benefit agent is an oil. For skin, preferred benefit agents include

- a) silicone oils, gums and modifications thereof such as linear and cyclic polydimethylsiloxanes; amino, alkyl alkylaryl and aryl silicone oils;
- b) fats and oils including natural fats and oils such as jojoba and beef tallow;
- c) waxes such as beeswax and lanolin;
- d) hydrocarbons such as petrolatum and mineral oil;
- e) higher fatty acids and higher fatty alcohols, both saturated and unsaturated, having a carbon chain length in the range C_{12} to C_{22} ;
- f) esters such as isopropyl myristate and isopropyl palmitate;
- g) essential oils such as evening primrose oil;
- h) lipids such as cholesterol, ceramides, sucrose esters and pseudo-ceramides as described in European Patent Specification No. 556 957;
- i) vitamins such as vitamin A and E, and vitamin alkyl esters, including those vitamin C alkyl esters;
- j) sunscreens such as octyl methoxyl cinnamate (Parsol MCX) and butyl methoxy benzoylmethane (Parsol 1789); and
- k) mixtures of any of the foregoing components.

For hair, the oil may take the form of a sunscreen, a styling or bodying agent or a conditioning oil. Among suitable

- 7 -

5 sunscreens and other benefit agents there may be mentioned:
the group of branched hydrocarbon materials of high molecular
weight referred to elsewhere as peralk(en)yl hydrocarbons.
These may be either in an organic solvent or directly
emulsified in the shampoo composition to give styling or
bodying effects. Polyisobutene is a preferred branched
hydrocarbon material. Also oil soluble sunscreens
partitioned into emulsified oil droplets. Among the oils
suitable for this purpose are phenyl silicones and among the
10 suitable sunscreens are, benzophenone compounds, dibenzoyl
methane derivatives and camphor derivatives. A preferred
sunscreen material is a UV absorber such as 2-ethyl hexyl
methoxy cinnamate sold under the trade name Parsol MCX by
Givaudan.

15 Silicone oil is a preferred conditioning oil for skin or
hair. The silicone may be in the form of a low viscosity oil
which may contain a high viscosity oil or gum in solution.
Alternatively the high viscosity material may be in the form
of an emulsion in water. The emulsion may be of high
20 viscosity oil or of a solution of gum in a lower viscosity
oil. The particle size of the oil phase may be anywhere in
the range from 30 nanometres to up to 20 microns average
size.

25 When the oil is a silicone it may be a polydimethylsiloxane
with an average particle size of less than 20 microns and
preferably less than 2 microns. Small particle size enables
a more uniform distribution of silicone conditioning agent
30 for the same concentration of silicone in the composition.
Advantageously a silicone with a viscosity in the range 1-20
million cst is used. The silicone can be cross-linked.

35 The personal washing composition may further comprise from
0.1 to 5 % of a suspending agent selected from polyacrylic


- 8 -

acids, cross linked polymers of acrylic acid, copolymers of acrylic acid with a hydrophobic monomer, copolymers of carboxylic acid- containing monomers and acrylic esters, cross-linked copolymers of acrylic acid and acrylate esters, heteropolysaccharide gums and crystalline long chain acyl derivatives. The long chain acyl derivative is desirably selected from ethylene glycol stearates, alkanolamides of fatty acids having from 16 to 22 carbon atoms and mixtures thereof. Ethylene glycol distearate and Polyethylene glycol 3 distearate are preferred long chain acyl derivatives. Polyacrylic acid is available commercially as Carbopol 420, Carbopol 488 or Carbopol 493. Polymers of acrylic acid cross-linked with a polyfunctional agent may also be used, they are available commercially as Carbopol 910, Carbopol 934, Carbopol 940, Carbopol 941 and Carbopol 980. An example of a suitable copolymer of a carboxylic acid containing a monomer and acrylic acid esters is Carbopol 1342. All Carbopol materials are available from Goodrich and Carbopol is a trade mark. The suspending agent is particularly preferred when silicone is present.

Suitable cross linked polymers of acrylic acid and acrylate esters are Pemulen TR1 or Pemulen TR2. A suitable heteropolysaccharide gum is xanthan gum, for example that available as Kelzan mu.

In skin washing compositions of the invention, the surface active agent can be selected from any known surfactant suitable for topical application to the human body. Mild surfactants, ie. surfactants which do not damage the stratum corneum, the outer layer of skin, are particularly preferred.


One preferred anionic surfactant is fatty acyl isethionate of formula:



- 9 -

where R is an alkyl or alkenyl group of 7 to 21 carbon atoms and M is a solubilising cation such as sodium, potassium, ammonium or substituted ammonium. Preferably at least three quarters of the RCO groups have 12 to 18 carbon atoms and may be derived from coconut, palm or a coconut/palm blend.

Another preferred anionic surfactant is alkyl ether sulphate of formula:



where R is an alkyl group of 8 to 22 carbon atoms, n ranges from 0.5 to 10 especially 1.5 to 8, and M is a solubilising cation as before.

Other possible anionic surfactants include alkyl glyceryl ether sulphate, sulphosuccinates, taurates, sarcosinates, sulphoacetates, alkyl phosphate, alkyl phosphate esters and acyl lactylate, alkyl glutamates and mixtures thereof.

Sulphosuccinates may be monoalkyl sulphosuccinates having the formula: $\text{R}^5\text{O}_2\text{CCH}_2\text{CH}(\text{SO}_3\text{M})\text{CO}_2\text{M}$; and amido-MEA sulphosuccinates of the formula: $\text{RCONCH}_2\text{CH}_2\text{O}_2\text{CCH}_2\text{CH}(\text{SO}_3\text{M})\text{CO}_2\text{M}$; wherein R^5 ranges from C_8 - C_{20} alkyl, preferably C_{12} - C_{15} alkyl and M is a solubilising cation.

Sarcosinates are generally indicated by the formula: $\text{R}^5\text{CON}(\text{CH}_3)\text{CH}_2\text{CO}_2\text{M}$, wherein R ranges from C_8 - C_{20} alkyl, preferably C_{12} - C_{15} alkyl and M is a solubilising cation.

Taurates are generally identified by the formula: $\text{R}^5\text{CONR}^6\text{CH}_2\text{CH}_2\text{SO}_3\text{M}$, wherein R^5 ranges from C_8 - C_{20} alkyl, preferably C_{12} - C_{15} alkyl, R^6 ranges from C_1 - C_4 alkyl, and M is a solubilising cation.

- 10 -


Harsh surfactants such as primary alkane sulphonate or alkyl benzene sulphonate will generally be avoided.

Suitable nonionic surface active agents include alkyl polysaccharides, lactobionamides, ethyleneglycol esters, glycerol monoethers, polyhydroxyamides (glucamide), primary and secondary alcohol ethoxylates, especially the C₈₋₂₀ aliphatic alcohols ethoxylated with an average of from 1 to 20 moles of ethylene oxide per mole of alcohol.

If the surface active agent comprises soap, the soap is preferably derived from materials with a C₈ to C₂₂ substantially saturated carbon chain and, preferably, is a potassium soap with a C₁₂ to C₁₈ carbon chain.

Mixtures of any of the foregoing surface active agents may also be used.

It is also preferable that the composition includes at least one cosurfactant agent with skin-mildness benefits. Suitable materials are zwitterionic detergents which have an alkyl or alkenyl group of 7 to 18 carbon atoms and comply with an overall structural formula



where R¹ is alkyl or alkenyl of 7 to 18 carbon atoms
R² and R³ are each independently alkyl, hydroxyalkyl or carboxyalkyl of 1 to 3 carbon atoms

m is 2 to 4


n is 0 or 1

- 11 -


X is alkylene of 1 to 3 carbon atoms optionally substituted with hydroxyl, and

Y is $-\text{CO}_2^-$ or $-\text{SO}_3^-$

5 Zwitterionic detergents within the above general formula include simple betaines of formula:



and amido betaines of formula:



where m is 2 or 3.


In both formulae R^1 , R^2 and R^3 are as defined previously. R^1 may, in particular, be a mixture of C_{12} and C_{14} alkyl groups derived from coconut so that at least half, preferably at least three quarters of the groups R^1 have 10 to 14 carbon atoms. R^2 and R^3 are preferably methyl.

A further possibility is a sulphobetaine of formula:




- 12 -

or



5

where m is 2 or 3, or variants of these in which
 $-(\text{CH}_2)_3\text{SO}_3^-$ is replaced by



10

R^1 , R^2 and R^3 in these formulae are as defined previously.

15

The surface active agent is preferably present in amount of
 from 2 to 40% by weight, and preferably from 5 to 30% by
 weight. The cosurfactant, is present, is preferably present
 at a level of 0.5 to 15% by weight.

20

The skin washing composition according to the invention may
 also include minor amounts of other ingredients such as
 antibacterial agents, foam boosters, pearlescers, perfumes,
 dyes, colouring agents, preservatives, thickeners, proteins,
 other polymers, phosphate esters and buffering agents.

25

Shampoo compositions of the invention contain anionic
 surfactant together with optional nonionic and amphoteric
 surfactant.

30

Suitable anionic surfactants are the alkyl sulphates, alkyl
 ether sulphates, alkaryl sulphonates, alkyl succinates, alkyl
 sulphosuccinates, N-alkoyl sarcosinates, alkyl phosphates,
 alkyl ether phosphates, alkyl ether carboxylates, and alpha-
 olefin sulphonates, especially their sodium, magnesium,
 ammonium and mono-, di- and triethanolamine salts. The alkyl
 groups generally contains from 8 to 18 carbon atoms and may
 be unsaturated. The alkyl ether sulphates, alkyl ether
 phosphates and alkyl ether carboxylates may contain from one

35

- 13 -

to 10 ethylene oxide or propylene oxide units per molecule, and preferably contain an average of 2 to 3 ethylene oxide units per molecule.

5 Further examples of suitable anionic surfactants include sodium oleyl succinate, ammonium lauryl sulphosuccinate, ammonium lauryl sulphate, sodium dodecylbenzene sulphonate, triethanolamine dodecylbenzene sulphonate, triethanolamine dodecylbenzene sulphonate and sodium N-lauryl sarcosinate.
10 The most preferred anionic surfactants are sodium lauryl sulphate, triethanolamine lauryl sulphate, triethanolamine monolauryl phosphate, sodium lauryl ether sulphate 1EO, 2EO and 3EO, ammonium lauryl sulphate and ammonium lauryl ether sulphate 1EO, 2EO and 3EO.

15 The nonionic surfactants suitable for use in the shampoo compositions of the invention include condensation products of aliphatic (C_8-C_{18}) primary or secondary linear or branched chain alcohols or phenols with alkylene oxides, usually
20 ethylene oxide and generally 6-30 EO.

Other suitable nonionics include mono or di alkyl alkanolamides or alkyl polyglucosides. Examples include coco mono or diethanolamide, coco mono isopropanolamide, and coco
25 di glucoside.

The amphoteric surfactants suitable for use in the composition of the invention may include alkyl amine oxides, alkyl betaines, alkyl amidopropyl betaines, alkyl
30 sulphobetaines, alkyl glycinate, alkyl carboxyglycinates, alkyl amphopropionates, alkyl amidopropyl hydroxysultaines, acyl taurates and acyl glutamates wherein the alkyl and acyl groups have from 8 to 18 carbon atoms. Examples include lauryl amine oxide, cocodimethyl sulphopropyl betaine and

- 14 -

preferably lauryl betaine, cocamidopropyl betaine and sodium cocamphopropionate.

5 The surfactants are present in the shampoo compositions of the invention in an amount of from 2 to 40% by weight, and preferably from 5 to 30% by weight.

10 The shampoo may also include minor amounts of other ingredients such as antibacterial agents, foam boosters, pearlescers, perfumes, dyes, colouring agents, preservatives, thickeners, proteins, other polymers, phosphate esters and buffering agents.

15 The invention will now be described, with reference to the following non-limiting examples:

[A] Skin Washing Compositions

Examples

20

In the examples:-

Coco amidopropyl betaine was Tergobetaine F ex Goldschmidt or Amonyl BA 380 ex Seppic.

25 Guar hydroxypropyl trimonium chloride was Jaguar C-13-S ex Meyhall.

Silicone oil emulsion was BC 89/138 ex Basildon.

Sodium cocoyl isethionate was either Jordapon CI ex PPG/Mazer or Hostapan SCI ex Hoechst.

30 Sodium lauryl ether sulphate was Genapol ZRO ex Hoechst.

Compositions according to the invention and comparative compositions were tested by the following method.

35 A number of tests were carried out by human volunteers. The experimental procedure employed was as follows:

- 15 -

The volunteers washed their forearms with a shower gel. The procedure involved wetting the arm and also the volunteer's free hand with warm water then using the free hand to lather the arm with 0.5 grams of the shower gel, next rinsing for 10 seconds while rubbing with the free hand and then drying the arm with a single pass with a paper towel.

10 minutes after drying the forearm a strip of adhesive tape is pressed onto the areas on the forearms keeping it in place for 30 seconds using a spring loaded device bearing on a rubber bung to press the tape onto the skin with a repeatable pressure of 85g.cm^{-2} . The adhesive tape employed was J-Lar Superclear (TM) tape having a width of 25mm. Two strips of tape are applied to each forearm in this way to adjacent areas of the skin.

In this test procedure silicone which has deposited on the skin is transferred to the tape along with some of the outer layer of the volunteer's skin.

The amounts of silicon and skin adhering to the tape are determined by means of X-ray fluorescence spectroscopy. The tape strips are placed in an X-ray fluorescence spectrometer with the adhesive side facing the beam of this machine. A mask is applied over the tape to define a standardised area in the middle of the tape which is exposed to the X-ray beam. The sample chamber of the machine is placed under vacuum before making measurements and the spectrometer is then used to measure the quantities of silicon and sulphur. The sulphur is representative of the amount of skin which has transferred to the tape. Results are presented in terms of the ratio of Si:S.

- 16 -

Example 1

In this example deposition of silicone from compositions containing a range of polymers according to the invention was compared with deposition from a composition containing a polymer which is a commercially available shower gel, namely, guar hydroxypropyl trimonium chloride (comparison).

The base formulation was:-

10

	%wt
Sodium lauryl ether sulphate (SLES)	13.00
Coco amidopropyl betaine (CAPB)	2.00
Silicone oil emulsion	5.00
15 Sorbic acid	0.37
Sodium citrate dihydrate	0.49
Sodium chloride	2.0
Citric acid	0.01
20 Water + minors	to 100

20

Polymers were added to the base formulation at a level of 0.1 %wt.

25

Each composition was prepared by forming a 1% dispersion of the polymer by adding it to water at ~50°C. SLES and CAPB were added to the excess water of the formulation with gentle stirring. Thereafter the silicone oil emulsion was added with stirring to the surfactant mixture. This was followed by the polymer dispersion and finally the minors.

30

- 17 -

Deposition of silicone was determined according to the procedure described above. The Si:S ratio for the comparison was normalised to 1 and the values for the compositions according to the invention expressed relative to the comparison.

The following results were obtained:-

Polymer	Si:S
Comparison	1
CPA 6	4.19
CPA 1	2.42
CPA 7	1.90

The results demonstrate the improved deposition obtained with the compositions according to the invention.

In a further set of experiment with the same base formulation but with different polymers the following results were obtained.

Polymer	Si:S
Comparison	1
CPA 9	1.83
CPA 10	2.10
CPA 5	2.81
CPA 8	1.72

- 18 -

These results also demonstrate the improved deposition obtained with the compositions according to the invention.

Example 2

In this example the variation of silicone deposition with the amount of polymer added to the base formulation was examined and compared with that from a composition containing no polymer.

The base formulation was:-

	%wt
Sodium lauryl ether sulphate	2.00
Coco amidopropyl betaine (CAPB)	8.00
Sodium cocoyl isethionate	5.00
Silicone oil emulsion	5.00
Water + minors	to 100

It was prepared by forming a premix of the cocoyl isethionate (25% dispersion) by adding it to water at 45°C. The SLES and isethionate premix were then added to the excess water of the formulation with gentle stirring, followed by the CAPB.

Thereafter the silicone oil emulsion was added with stirring. A 1% dispersion of the polymer was prepared by adding it to water at ~50°C. This was then added to the surfactant/silicone mixture to the required level followed by the minors.

Deposition of silicone was determined according to the procedure described above.

- 19 -

The following results were obtained:-

Polymer	%wt	Si:S
Comparison	0	0.41
CPA 5	0.05	1.02
	0.1	2.43
	0.2	3.28
Comparison	0	0.36
CPA 6	0.05	2.95
	0.1	4.28
	0.2	4.16

The results demonstrate that silicone deposition increases as the amount of polymer present in the composition increases.

[B] Shampoo Compositions

Test of Conventional Cationic Polymers

Many of the commercially available cationic polymers designed for use in cosmetics show no ability to deposit silicone on to hair during the course of the hair washing/rinsing cycle. Table 1 details the performance of a range of cationic polymers promoted by their manufacturers as suitable for use in shampoo applications. The polymers were tested as silicone deposition and retention aids in one of the shampoo formulations (A,B,C or D) given below in Table 3.

- 20 -

Table 1

Eg	Polymer	shampoo	silicone retention on hair ppm	% deposition	polymer Mw
A	C13S	C	1280	26	250 000
B	C13S	A	983	20	250 000
C	C15	A	142	3	<100 000
D	C13S	D	1828	25	250 000
E	JR400	A	0	-	400 000
F	JR30M	A	0	-	400 000
G	JR400	B	661	14	400 000
H	JR400	C	0	-	400 000
I	FC370	A	0	-	100 000
J	Quat- PVA	A	0	-	125 000
K	550	A	0	-	700 000
L	-	A	0	-	-
M	-	C	0	-	-

A figure of zero for silicone retention indicates that the amount detected was negligible and could not accurately be measured.

C13S is JAGUAR (trade mark) C13S, a cationic guar derivative ex Meyhall

- 21 -

C15 is JAGUAR C15, also a cationic guar derivative ex Meyhall

JR400 is POLYMER JR400, a polysaccharide derivative ex Union Carbide

5 JR30M is POLYMER JR30M, a polysaccharide derivative ex Union Carbide

FC370 is Luviquat FC 370 (trade mark), ex BASF

Quat-pva is a copolymer prepared by reacting glycidyltrimethylammonium chloride with a commercial PVA;

10 Mowiol 40-88, ex Hoeschst. The molecular weight (Mw) is 127,000 (supplier's data). The final charge density (from NMR) is 1.2 meq/g.

550 is Merquat 550 ex Croxton and Garry.

15

Of these examples, only example K is polyacrylamide based. The value given for polymer Mw in the table is suppliers data for all examples except Example K. According to Croxton + Garry, Merquat 550 has a weight average molecular weight of 2.8 million, however, when measured using our Intrinsic viscosity method a value for Mw of 700 000 is obtained.

20

All retention figures were obtained direct from hair switches washed twice for 30s, and rinsed twice for 30s, in running tap water. Shampoo application was at the level of 0.12g/g hair. Silicone levels were determined from X-Ray Fluorescence count rates by comparison with known standards.

25

Silicone retention efficiency from formulations containing these low Mw polymers does not exceed 25%.

30

Examples 3-8 High Molecular Weight Cationic Polyacrylamides

Using high molecular weight polyacrylamides (Mw>3,000,000) gives previously unattainable levels of silicone retention,

35

- 22 -

typically far in excess of 50% efficiency. The results are shown in Table 2.

Table 2

Example	Polymer	Shampoo	Silicone retention ppm	Efficiency %	Mol wt
3	CPA1	D	>5000	>70	8 000 000
4	CPA1	A	2060	43	8 000 000
5	CPA2	A	0	-	400 000
6	CPA3	C	63	63	6 000 000
7	CPA4	C	130	3	220 000
8	'703	A	>>2500	>>55	>5 000 000

Thus, although most commercially available cationic polymers with Mw below 1,000,000 intended for use in cosmetic products show little or no activity as deposition aids from conventional shampoo formulations, use of very high Mw (> 3,000,000) cationic polyacrylamides in shampoo formulations gives surprisingly increased levels of silicone retention.

- 23 -

Table 3SHAMPOO A:

5 16 SLES 2EO
 2 Lauryl Betaine
 2.25 Ethylene glycol distearate
 4 BY22-026 (50% silicone emulsion) ex Toray silicone
10 0.1 Deposition Polymer as specified

SHAMPOO B:

 8 SLES 3EO
 4 cocoamido propylbetaine
15 1.5 NaCl
 4 BY22-026
 0.3 Deposition Polymer as specified

SHAMPOO C:

20 12 SLES 2EO
 2 BY22-026
 0.1 Deposition Polymer as specified

25 SHAMPOO D:

 16 SLES 2EO
 2 cocoamido propylbetaine
 1.5 ethylene glycol distearate
30 0.5 Potassium sorbate
 0.25 Citric Acid
 4.8 X2-1766 (60% Silicone emulsion) ex Dow Corning
 0.3 NaCl
 0.1 Deposition Polymer as specified

35

- 24 -

Comparative Examples N and O

Because we believed that the higher charge density Jaguar C17 might outperform the Jaguar C13S tested in Examples A, B and D, we made a comparison between C17 and other polymers in shampoo A as follows:

Example N: Shampoo A with Jaguar C13S 983 ppm 20% efficiency

Example O: Shampoo A with Jaguar C171 413 ppm 29% efficiency

The polyacrylamides with high molecular weight (Examples 1, 2, 4 and 6) clearly outperform Jaguar C17.

The silicone level in these shampoos was 2% by weight.

Example 9 and Comparative Examples P and Q

To show that the deposition polymers according to the invention can give benefit when added to a commercial shampoo we tested "Wash and Go" Dry Sensitive, (Comparative Example P believed to be without deposition polymer) and "Wash and Go" Extra Conditioning (Comparative Example Q, believed to be with Jaguar C17) against the "Dry Sensitive" formulation with 0.2% CPA1 added, (Example 7). CPA1 is a copolymer of acrylamide and N,N,N-trimethylaminopropylacrylamide. "Wash and Go" is a range of 2 in 1 shampoos sold by Procter & Gamble based on an alkyl sulphate and ether sulphate anionic surfactant mixture. The increased deposition from use of the deposition polymer according to the invention is readily apparent.

Deposition of silicone (ppm silicone) was measured by X-Ray Fluorescence:

- 25 -

Example P 562 +/- 289 ppm

Example Q 468 +/- 103 ppm

5 Example 7 1630 +/- 635 ppm


All figures are an average from 5 hair samples. It can be seen that addition of the Polymer according to the invention trebles the efficiency of silicone deposition.

10

- 26 -

CLAIMS

1. A personal washing composition comprising:-
- a) a surface active agent selected from the group consisting of anionic, nonionic, zwitterionic and cationic surfactants, soap and mixtures thereof;
 - b) water;
 - c) a non-volatile insoluble benefit agent dispersed in the composition; and
 - d) from 0.001 to 1% by weight of a deposition polymer which is a cationic copolymer wherein the charge density of the copolymer is in the range 0.0001 to 0.005 eq/g; and the average molecular weight of the copolymer is more than 2×10^6 daltons.
2. A composition according to claim 1 in which the amount of deposition polymer lies in the range 0.05 to 0.2% by weight.
3. A composition according to claim 1 or 2 in which the cationic copolymer is a copolymer of acrylamide and a cationic monomer having the formula:



where: T is -O- or -C-



R is H or CH_3

and R' is $-\text{NH}-(\text{CH}_2)_n-\text{N}^+(\text{CH}_3)_3 \text{X}^-$ or $-\text{O}-(\text{CH}_2)_n-\text{N}^+(\text{CH}_3)_3 \text{X}^-$

- 27 -

in which n is an integer from 1 to 4 and X is selected from Cl, Br, I and CH_3SO_3 .

4. A composition according to any preceding claim in which the charge density of the deposition polymer lies in the range 0.0008 to 0.0025 eq/g.
5. A composition according to any preceding claim in which the benefit agent is selected from the group consisting of silicone oils and gums; fats; oils; waxes; hydrocarbons; higher fatty acids; higher fatty alcohols; esters; essential oils; lipids; vitamins; sunscreens; and mixtures thereof.
6. A composition according to claim 5 in which the benefit agent is a silicone oil.
7. A composition according to claim 6 in which the composition further comprises from 0.1 to 5 % of a suspending agent for the silicone oil selected from the group consisting of polyacrylic acids; cross linked polymers of acrylic acid; copolymers of acrylic acid with a hydrophobic monomer; copolymers of carboxylic acid- containing monomers and acrylic esters; cross-linked copolymers of acrylic acid and acrylate esters; heteropolysaccharide gums; crystalline long chain acyl derivatives; fatty acid monoglyceride polyglycol ethers; propylene glycol and propylene glycol oleate; and mixtures thereof.
8. A composition according to any preceding claim which is a shampoo composition and in which the surface active agent is an anionic surfactant.

- 28 -

9. A shampoo composition according to claim 8 in which the anionic surfactant is selected from the group consisting of sodium lauryl sulphate, triethanolamine lauryl sulphate, triethanolamine monolauryl phosphate, sodium lauryl ether sulphate 1EO, 2EO and 3EO, ammonium lauryl sulphate and ammonium lauryl ether sulphate 1EO, 2EO and 3EO and mixtures thereof.
10. A shampoo composition according to claim 8 or 9 in which the benefit agent is selected from the group consisting of sunscreens, styling or bodying agents and conditioning oils.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 95/00582

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K7/48 A61K7/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,92 10162 (THE PROCTER & GAMBLE COMPANY) 25 June 1992 see the whole document ---	1-10
X	WO,A,93 08787 (THE PROCTER & GAMBLE COMPANY) 13 May 1993 see the whole document ---	1-10
X	US,A,4 299 817 (HANNAN ROY B.) 10 November 1981 see the whole document ---	1-10
X	EP,A,0 524 434 (HELENE CUSTIS INC.) 27 January 1993 see the whole document ---	1-10
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * & * document member of the same patent family

Date of the actual completion of the international search

13 July 1995

Date of mailing of the international search report

26 / 07 / 95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo.nl,
Fax: (+31-70) 340-3016

Authorized officer

Couckuyt, P

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 95/00582

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP,A,0 093 601 (UNILEVER PLC) 9 November 1983 see the whole document ---	1-5,7-10
X	WO,A,93 07903 (RICHARDSON-VICKS INC.) 29 April 1993 see the whole document ---	1,3,5,6, 10
P,X	FR,A,2 698 004 (L'OREAL) 20 May 1994 see the whole document ---	1-7,9
X	EP,A,0 219 830 (DOW CORNING CORPORATION) 29 April 1987 see the whole document ---	1,5-7
P,X	WO,A,94 21224 (L'OREAL) 29 September 1994 see the whole document ---	1-10
E	WO,A,95 09599 (UNILEVER PLC) 13 April 1995 see the whole document -----	1-10

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 95/00582

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9210162	25-06-92	AT-T- 111727	15-10-94
		AU-A- 9117791	08-07-92
		CA-A- 2097836	06-06-92
		CN-A- 1062837	22-07-92
		DE-D- 69104224	27-10-94
		DE-T- 69104224	09-03-95
		EP-A- 0560919	22-09-93
		ES-T- 2060467	16-11-94
		FI-A- 932565	04-06-93
		HU-A- 64686	28-02-94
		JP-T- 6503574	21-04-94
		NZ-A- 240856	24-02-95
WO-A-9308787	13-05-93	AU-A- 2927292	07-06-93
		BR-A- 9206689	02-05-95
		CA-A- 2120492	13-05-93
		CZ-A- 9401025	19-10-94
		EP-A- 0610407	17-08-94
		FI-A- 941969	28-04-94
		HU-A- 67994	14-03-95
		JP-T- 7500609	19-01-95
		NO-A- 941535	28-06-94
		PT-A- 101011	28-02-94
US-A-4299817	10-11-81	NONE	
EP-A-524434	27-01-93	US-A- 5221530	22-06-93
		AU-B- 652257	18-08-94
		AU-A- 1847292	07-01-93
		CA-A- 2070299	25-12-92
		JP-A- 5186317	27-07-93
		US-A- 5417965	23-05-95
EP-A-93601	09-11-83	AU-B- 546461	05-09-85
		AU-A- 1398983	03-11-83
		CA-A- 1203141	15-04-86
		DE-A- 3374018	12-11-87
		GB-A, B 2122214	11-01-84
		JP-C- 1319715	29-05-86
		JP-A- 58196300	15-11-83

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 95/00582

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-93601		JP-B- 60046158 US-A- 5037818	14-10-85 06-08-91
WO-A-9307903	29-04-93	AU-A- 2863992 CA-A- 2117265 CZ-A- 9400901 EP-A- 0608322 FI-A- 941770 HU-A- 67046 JP-T- 7500594 NO-A- 941317 PT-A- 100964	21-05-93 29-04-93 13-07-94 03-08-94 15-04-94 30-01-95 19-01-95 16-06-94 30-11-93
FR-A-2698004	20-05-94	EP-A- 0603019 JP-A- 6211626 US-A- 5368850	22-06-94 02-08-94 29-11-94
EP-A-219830	29-04-87	US-A- 4744978 AU-B- 586709 AU-A- 6433686 CA-A- 1270443 DE-A- 3684088 JP-A- 62103007 KR-B- 9401000	17-05-88 20-07-89 30-04-87 19-06-90 09-04-92 13-05-87 08-02-94
WO-A-9421224	29-09-94	FR-A- 2702653 AU-B- 6260694	23-09-94 11-10-94
WO-A-9509599	13-04-95	AU-B- 7810594	01-05-95